



## Review Article

# Syndrome resembling Kawasaki disease in COVID-19 asymptomatic children

Suriya Rehman<sup>a,\*</sup>, Tariq Majeed<sup>b</sup>, Mohammad Azam Ansari<sup>a</sup>, Ebtessam A. Al-Suhaimi<sup>c,\*</sup>

<sup>a</sup> Department of Epidemic Diseases Research, Institute of Research and Medical Consultations (IRMC), Imam Abdulrahman Bin Faisal University, 31441 Dammam, Saudi Arabia

<sup>b</sup> Department of General Pediatric, Maternity and Children Hospital, Dammam, Saudi Arabia

<sup>c</sup> Department of Biology, College of Science and Institute of Research and Medical Consultations (IRMC), Imam Abdulrahman Bin Faisal University, 31441 Dammam, Saudi Arabia

## ARTICLE INFO

## Article history:

Received 16 June 2020

Received in revised form 7 August 2020

Accepted 13 August 2020

## Keywords:

COVID-19

Children

Multisystem inflammatory syndrome

Kawasaki disease

## ABSTRACT

The current knowledge about the COVID-19 (Coronavirus Disease-2019) pandemic is still limited and is unravelling with the passing days, especially clinical data, and research in pediatric age group. Recently, there is a new and crucial development reported recently among the COVID-19 asymptomatic children, a novel syndrome affecting asymptomatic COVID-19 children, presenting as a hyperinflammatory syndrome which is like Kawasaki disease shock syndrome. The purpose of this correspondence is to discuss some important findings of the syndrome for the better understanding of the disease.

© 2020 The Authors. Published by Elsevier Ltd on behalf of King Saud Bin Abdulaziz University for Health Sciences. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Contents

Conclusion.....	1831
Funding .....	1831
Competing interests.....	1831
Ethical approval.....	1831
References.....	1831

Until now, children are considered as the population that is minimally affected by COVID-19 [1]. However, after the start of covid-19 pandemic, clinicians in many parts of the world have recognized a new syndrome resembling Kawasaki disease (vasculitis) in pediatric patients [2]. The syndrome has been named as pediatric multisystem inflammatory syndrome (MIS-C) and is characterized by high fever rash hypotension, gastrointestinal symptoms, and organ dysfunction. All patients finally developed warm vasoplegic shock unresponsive to volume resuscitation. Other features besides fever and rash included small pleural pericardial and ascitic effusions, suggestive of a widespread inflammatory process [3]. Surprisingly, respiratory symptoms do not seem to be common. Although this syndrome resembles Kawasaki disease in many

aspects, but the classical symptoms of Kawasaki disease, like bilateral conjunctival injection, strawberry tongue and rash, are not always present [4]. According to study published in Lancet, among the eight kids between the age of 4 and 14 years, three needed mechanical ventilation, and one died. The investigation showed either positive RTPCR (reverse transcriptase polymerase chain reaction) or positive antibodies to SARS-COV-2 (Severe Acute Respiratory Syndrome Coronavirus), which suggests that this syndrome occurs in the convalescent phase of the disease. The highest risk of this syndrome has been found to be between the ages 5 and 14 years. However, the risk for this syndrome remains very small [3].

Laboratory investigation revealed that MIC-S has far more elevated CRP (C-reactive protein), D dimer ferritin, elevated cardiac enzymes, and low lymphocyte count in CBC (complete blood count), compared to Kawasaki syndrome. The complications of this syndrome include multi organ failure, thrombosis, ischemic events, and coronary aneurysms. ECG (electrocardiogram) were found to be nonspecific and echocardiograph showed eco bright coronary arteries which progressed to coronary aneurysm in one patient [5].

\* Corresponding authors.

E-mail addresses: [surrehman@iau.edu.sa](mailto:surrehman@iau.edu.sa), [suriyamir@gmail.com](mailto:suriyamir@gmail.com) (S. Rehman), [ealsuhaimi@iau.edu.sa](mailto:ealsuhaimi@iau.edu.sa) (E.A. Al-Suhaimi).

Regarding the treatment all patients were treated with intravenous immunoglobulins and aspirin, however aspirin should be withheld as it can lead to bleeding episodes. All the children were discharged from ICU (intensive care unit) after 4–6 days. All children have been given follow up with cardiologist to rule out coronary abnormalities caused by the virus.

In one of the case studies done in United Kingdom, 8 patients presented with fever, conjunctivitis, extremity pain, edema, and diarrhea and all the patients had fluid refractory shock requiring vasopressors. Surprisingly, there were minimal respiratory symptoms, 50% of these patients had a family exposure to SARS-CoV-2, but RT-PCR was found to be negative. Majority of patients had some cardiac dysfunction and increased inflammatory markers and troponin [6]. In Bergamo, Italy, a comparative study was done between 19 patients who presented with Kawasaki like illness before the pandemic, to ten patients who presented with Kawasaki like illness after the start of pandemic. Those patients who presented after the start of pandemic had an average age of 7.5 years, while for the other group, the average age was 3 years and 50% of the patients had incomplete features of Kawasaki disease. Non-Kawasaki signs and symptoms included diarrhea, meningeal signs, and symptoms of hypoperfusion. Most of the patients had neutrophilia hyponatremia, hypertriglyceridemia, high ESR (erythrocyte sedimentation rate), CRP, ferritin, and pro-BNP (brain natriuretic peptide). Only two patients had positive swab for SARS-CoV-2. These two patients and six other patients have positive serologies, whereas, 50% of 10 patients had positive x-ray (radiograph) and 60% had abnormal echocardiography [7].

In another study among households in Shenzhen, China, children below the age of 10 years were found positive for COVID-19, as adults, however, they were found to have less severe symptoms [8]. Likewise, other reports from various countries including South Korea, Italy, and Iceland, have reported less rates of infection among children than the adults on the basis of mass testing [9]. Some preliminary data of almost 300 infected children suggested that lower concentration of cytokines is being produced by children compared to the affected adults. These cytokines are the proteins secreted by the body's immune system and is constant with Kawasaki disease as a disease related with the dysfunction of acquired immune system. Cytokines slows down the function of heart *via* activation of direct immune stimulation associated with myocytolysis [10].

There is enough data that suggests the resemblance of the symptoms of this infection and Kawasaki disease. However, differences are also seen, like age appears to be higher than the classical Kawasaki disease which is usually 3. Additionally, it has been found African children are more to MIS-C while Kawasaki is more common in children of Asian ancestry, indicating a racial predisposition. Another striking point is the gastrointestinal symptoms, SARS-CoV-2 have the potential to infect enterocytes *via* ACE2 (angiotensin converting enzyme 2) receptors being expressed in gastric epithelia which can lead to diarrhea in approximately 1/3 infected adults. On the contrary, occurrence of diarrhea in a post COVID-19 disease is likely not associated directly with infection. In typical Kawasaki disease, emesis is less common than in COVID-19 linked syndrome (Riphagen et al. [3]) [12]. The syndrome associated with covid-19 seems to present more gastrointestinal symptoms like vomiting, diarrhea and abdominal pain, compared to classical Kawasaki diseases. Also, in various studies it has been found that MIS-C presents with low platelet count while Kawasaki presents with increased platelet count. It cannot be excluded that the MIS-C present different unrelated Kawasaki disease, atypical Kawasaki disease, toxic shock syndrome and myocarditis, which are different entities with different mechanisms [13]. For instance, a study on 9 patients suggested inverse correlation with Kawasaki disease as per the age. Another study from New York have reported the immune response is linked to the gene abnormalities [14].

Interestingly, it seems that this syndrome might not be specific to this virus, it may be that other viruses can also activate this hyperinflammation in children. There may be factors like genetic predisposition which may make some children susceptible to this syndrome [15]. Centers for Disease Control and Prevention (CDC), Royal college of Physicians for child health (RCPCH) and World Health Organization (WHO) have published few case definitions for this syndrome, which are as follows: (A) patient aged <21 years having fever (1) laboratory finding of inflammation, (2) symptoms of clinically severe illness which require hospital admission, with involvement of multiorgan system ( $\geq 2$ ) (respiratory, renal, hematologic, cardiac, gastrointestinal, neurological and dermatologic); (B) No alternative credible diagnoses; (C) currently or recently tested positive by RT-PCR for SARS-CoV-2 infection, or by serology, or antigen test; or exposure to COVID-19 within one month, before the appearance of symptoms, (1) Pyrexia  $>38.0^{\circ}\text{C}$  for more than 24 h, or record of subjective fever which lasts for more than 24 h, (2) Others includes, but not limited to the following criteria: an elevated acute phase reactants (CRP, ESR, procalcitonin, ferritin), fibrinogen, lactic acid dehydrogenase (LDH), D-dimer or interleukin 6 (IL-6), elevated neutrophils, reduced lymphocytes, and low albumin. Additionally, some patients may have complete criteria for Kawasaki disease others may not, but needs to be notified if they fulfil the case definition for MIS-C. Also, report the MIS-C in any pediatric mortality with positive SARS-CoV-2 infection [16,17,15].

## Conclusion

Therefore, our conclusion is that this may be a novel syndrome affecting asymptomatic COVID-19 children, presenting as a hyperinflammatory syndrome which is like Kawasaki disease shock syndrome. The relation between SARS-CoV-2 and MIS-C in children will continue to be elucidated by further research.

## Funding

No funding Sources.

## Competing interests

None declared.

## Ethical approval

Not required.

## References

- [1] Rehman S, Majeed T, Azam Ansari M, Ali U, Sabit H, Al-Suhaimi EA. Current scenario of COVID-19 in pediatric age group and physiology of immune and Thymus response. Saudi J Biol Sci 2020;(xxxx), <http://dx.doi.org/10.1016/j.sjbs.2020.05.024>.
- [2] Rauf A, Vijayan A, John ST, Krishnan R, Latheef A. Multisystem inflammatory syndrome with features of atypical Kawasaki disease during COVID-19 pandemic. Indian J Pediatr 2020;2–4, <http://dx.doi.org/10.1007/s12098-020-03357-1>.
- [3] Riphagen S, Gomez X, Gonzalez-Martinez C, Wilkinson N, Theocharis P. Hyperinflammatory shock in children during COVID-19 pandemic. Lancet (London, England) 2020;395(no. 10237):1607–8, [http://dx.doi.org/10.1016/S0140-6736\(20\)31094-1](http://dx.doi.org/10.1016/S0140-6736(20)31094-1).
- [4] Ma L, Zhang YY, Yu HG. Clinical manifestations of Kawasaki disease shock syndrome. Clin Pediatr (Phila) 2018, <http://dx.doi.org/10.1177/0009922817729483>.
- [5] Liu PP, Blet A, Smyth D, Li H. The science underlying COVID-19: implications for the cardiovascular system. Circulation 2020, <http://dx.doi.org/10.1161/circulationaha.120.047549>.
- [6] Jones VG, et al. COVID-19 and Kawasaki disease: novel virus and novel case. Hosp Pediatr 2020;(April):27–8, <http://dx.doi.org/10.1542/hpeds.2020-0123>.
- [7] Verdoni L, et al. An outbreak of severe Kawasaki-like disease at the Italian epicentre of the SARS-CoV-2 epidemic: an observational cohort study. Lancet 2020, [http://dx.doi.org/10.1016/s0140-6736\(20\)31103-x](http://dx.doi.org/10.1016/s0140-6736(20)31103-x).

- [8] Bi Q, et al. Epidemiology and transmission of COVID-19 in Shenzhen China: analysis of 391 cases and 1,286 of their close contacts. medRxiv 2020, <http://dx.doi.org/10.1101/2020.03.03.20028423>.
- [9] Lipshultz Steven E, Barach P. Rethinking COVID-19 in children: lessons learned from pediatric viral and inflammatory cardiovascular diseases. *Prog Pediatr Cardiol* 2020.
- [10] Zhang MM, Shi L, Li XH, Lin Y, Liu Y. Clinical analysis of Kawasaki disease shock syndrome. *Chin Med J* 2017, <http://dx.doi.org/10.4103/0366-6999.219151>.
- [12] Lamers MM, et al. SARS-CoV-2 productively infects human gut enterocytes. *Science* (80-.) 2020, <http://dx.doi.org/10.1126/science.abc1669>.
- [13] Scuccimarri R. Kawasaki disease. *Pediatr Clin North Am* 2012, <http://dx.doi.org/10.1016/j.pcl.2012.03.009>.
- [14] Reuters. Three U.S. Children With COVID-19 Have Rare Inflammatory Syndrome. *New York Time*; 2020, 04 28.
- [15] Care E, no. June EXTRA! COVID-19: The Impact on Pediatric; 2020.
- [16] Scientific brief. Multisystem Inflammatory Syndrome in Children and Adolescents with COVID-19, World Heal. Organ, [www.who.int](http://www.who.int).
- [17] Emergency.cdc.gov. Multisystem inflammatory syndrome in children (MIS-C) associated with coronavirus disease 2019 (COVID-19). *Centers Dis Control Prev* 2020;(May).